

# Constraints on N<sub>2</sub>O budget changes since pre-industrial time from new firn air and ice core isotope measurements

S. Bernard<sup>1</sup>, T. Röckmann<sup>2,\*</sup>, J. Kaiser<sup>2,3</sup>, J.-M. Barnola<sup>1</sup>, H. Fischer<sup>4</sup>, T. Blunier<sup>5</sup>, and J. Chappellaz<sup>1</sup>

<sup>1</sup>Laboratoire de Glaciologie et Géophysique de l'Environnement (CNRS-UJF), St Martin d'Hères, France

<sup>2</sup>Max-Planck Institute for Nuclear Physics, Heidelberg, Germany

<sup>3</sup>Department of Geosciences, Princeton University, Princeton, New Jersey, USA

<sup>4</sup>Alfred-Wegener Institute for Polar and Marine Research, Bremerhaven, Germany

<sup>5</sup>Climate and Environmental Physics Institute, University of Bern, Bern, Switzerland

<sup>\*</sup>now at: Institute for Marine and Atmospheric Research Utrecht, Utrecht University, Utrecht, The Netherlands

Received: 20 April 2005 - Accepted: 8 June 2005 - Published: 29 August 2005

Correspondence to: T. Röckmann (t.roeckmann@phys.uu.nl)

© 2005 Author(s). This work is licensed under a Creative Commons License.

7547

# Abstract

A historical record of changes in the  $N_2O$  isotope composition is important for a better understanding of the global  $N_2O$  atmospheric budget. Here we have combined measurements of trapped gases in the firn and in ice cores of one Arctic site (North GReen-

- <sup>5</sup> land Ice core Project NGRIP) and one Antarctic site (Berkner Island). We have performed measurements of the <sup>18</sup>O and position dependent <sup>15</sup>N isotopic composition of N<sub>2</sub>O. By comparing these data to simulations carried out with a firn air diffusion model, we have reconstructed the temporal evolution of the N<sub>2</sub>O isotope signatures since preindustrial times. The decrease observed for all signatures is consistent from one pole
- to the other. Results obtained from the air occluded in the ice suggest a decrease of about -2.8%, -2.4%, -3.2% and -1.6% for  $\delta^{15}$ N,  ${}^{1}\delta^{15}$ N,  ${}^{2}\delta^{15}$ N and  $\delta^{18}$ O, respectively, since 1700 AD. Firn air data imply a decrease of about -1.1%, -1.2%, -1.0% and -0.6% for  $\delta^{15}$ N,  ${}^{1}\delta^{15}$ N,  ${}^{2}\delta^{15}$ N and  $\delta^{18}$ O, respectively, since 1970 AD. These results imply consistent trends from firn and ice measurements for  $\delta^{15}$ N and  $\delta^{18}$ O. The
- trends for the intramolecular distribution of <sup>15</sup>N are less well constrained than the bulk <sup>15</sup>N trends because of the larger experimental error for the position dependent <sup>15</sup>N measurements. The decrease in the heavy isotope content of atmospheric N<sub>2</sub>O can be explained by the increasing importance of agriculture for the present atmospheric N<sub>2</sub>O budget.

## 20 1 Introduction

Over the past 200 years the mixing ratio of atmospheric  $N_2O$  has increased from ~270 nmol mol<sup>-1</sup> (Flückiger et al., 1999) to the value of 319 nmol mol<sup>-1</sup> in 2004. It is currently rising at a rate of (0.25±0.05)% per year (IPCC, 2001). Increasing  $N_2O$  levels are of concern for two reasons. Firstly,  $N_2O$  is a potent greenhouse gas and

<sup>25</sup> contributes a significant portion to anthropogenic climate change, even though it is more than 1100 times less abundant than carbon dioxide. This is due to the fact that the Global Warming Potential per molecule N<sub>2</sub>O over a 100-yr time horizon is about 300 times greater than for CO<sub>2</sub> (IPCC, 2001). Secondly, N<sub>2</sub>O plays an important role in ozone chemistry, because it is the main source of NO<sub>x</sub> in the stratosphere (Crutzen, 1970; Minschwaner et al., 1993). N<sub>2</sub>O is mainly produced by microbial nitrification and

- <sup>5</sup> denitrification processes in soils and water. Minor contributions come from biomass burning, industry, combustion in vehicles and power plants. The main sink of N<sub>2</sub>O is UV photolysis in the stratosphere (90%) with additional contributions from the reaction with electronically excited oxygen atoms (10%). Although precise estimates of the global N<sub>2</sub>O source can be deduced from the global rate of increase and the sink terms (Minashuanar et al. 1002) the centribution of individual equate terms remains peakly.
- (Minschwaner et al., 1993), the contribution of individual source terms remains poorly constrained.

A way to constrain the  $N_2O$  budget is to monitor its isotopic composition in the atmosphere (Dore et al., 1998; Kaiser et al., 2003a; Kim and Craig, 1993; Naqvi et al., 1998; Pérez et al., 2000, 2001; Röckmann et al., 2003a, 2001a, b; Sowers et al.,

- $^{15}$  2003, 2002; Yoshida and Toyoda, 2000). Because the different sources and sinks of  $N_2O$  have characteristic isotope signatures, the cause of the increase of atmospheric  $N_2O$  since pre-industrial times can be determined from the simultaneous evolution of its isotopic composition. Using a simple one box model accounting for the natural and anthropogenic sources and sinks of  $N_2O$ , Rahn and Wahlen (2000) predicted an iso-
- <sup>20</sup> topic difference between the pre-industrial and modern troposphere in the range -1.6 to -1.9‰ for  $\delta^{15}$ N and -1.8 to -2.4‰ for  $\delta^{18}$ O. Similarly, Pérez et al. (2001) proposed that the tropospheric  $\delta^{15}$ N of N<sub>2</sub>O should have dropped by -2.2 to -3‰ over the last two centuries, assuming that one half of the increase of N<sub>2</sub>O emissions is due to the use of synthetic N fertilizers.
- Air trapped in Arctic and Antarctic firn and ice provides a way to quantify the magnitude of the atmospheric N<sub>2</sub>O isotope changes. From analyses of Antarctic firn air, Röckmann et al. (2003a) extrapolated a decrease since the pre-industrial era, of -2%and -1.2% for  $\delta^{15}N$  and  $\delta^{18}O$ , respectively, whereas Sowers et al. (2002), using both firn air and ice core samples, suggested a decrease of -1.9% and -2.9% for  $\delta^{15}N$

7549

and  $\delta^{18}$ O, respectively.

Here, we reconstruct the isotopic composition of atmospheric  $N_2O$  since preindustrial times, using air from ice cores and firn, from an Arctic (North GRIP) and an Antarctic (Berkner Island) site. A firn-air diffusion model (Rommelaere et al., 1997) is used to take into account physical processes occurring in the firn column and to test

atmospheric scenarios of N<sub>2</sub>O isotope signatures against the measurements.

# 2 Experimental procedure

In the framework of the European Project CRYOSTAT (CRYospheric STudies of Atmospheric Trends of stratospherically and radiatively important gases), 15 samples of

- firn air were obtained from North GRIP in June 2001 (Greenland, 75° N 42° W) and from Berkner Island in January 2003 (Antarctica, 79° S 45° W). NGRIP (Berkner Island) has a mean annual temperature of  $-31^{\circ}$ C ( $-26^{\circ}$ C) and a surface accumulation rate of 20 cm (13 cm) water equivalent per year; its firn-ice transition is located at ~78 m (~64 m) and the mean N<sub>2</sub>O age at this depth is 60 years (90 years).
- The air sampling procedure is described in Bräunlich et al. (2001). Briefly, air samples were obtained by incrementally drilling into the firn with a standard electromechanical drill. Drilling was interrupted at each prescribed depth and a "bladder" made of natural rubber was lowered to the bottom of the hole and inflated, thereby sealing the hole. Firn air was then pumped, dried using magnesium perchlorate (MgClO<sub>4</sub>) and Drifte (achieve and whether the stand die Destelle Oliver Oliver, and
- Drierite (calcium sulfate) and ultimately stored in Restek Silcocans (3 liter, 3 bar) and Scott Marin cylinders (5 liter, 120 bar). In addition to the firn air, two ice samples from NGRIP and six ice samples from

Berkner Island were analyzed. A new technique to extract the air enclosed in the ice bubbles was used. A sample of ~1 kg is inserted into a glass container sealed with

an indium wire. After evacuation, the sample is melted under vacuum with continuous transfer of the released gases into an electropolished stainless steel cryotrap of 14 cm<sup>3</sup> held at liquid helium temperature. Two coiled glass traps and one stainless steel trap

between the container and the cryotrap are maintained at  $\sim -50^{\circ}$ C and  $-80^{\circ}$ C, respectively, in order to trap water vapor. The efficiency of the system was tested as following. To assure the absence of leaks in the extraction and transfer system, three artificial bubble-free ice samples of  $\sim 1$  kg each were melted and "transferred" into the cryotrap.

- <sup>5</sup> No detectable N<sub>2</sub>O or permanent gas could be detected. Next, to investigate possible isotopic fractionation, three ~1 kg bubble-free ice samples were processed together with ~100 cm<sup>3</sup> of an atmospheric standard having known N<sub>2</sub>O isotopic composition. Table 1 shows that overall no significant isotopic fractionation was observed between N<sub>2</sub>O after extraction and the standard gas measured on the same day straight from its
- <sup>10</sup> cylinder, although  $\delta^{18}$ O may be slightly enriched. Based on these tests, we chose to apply no extraction correction to the ice-core results.

A potential source of error is the dissolution of N<sub>2</sub>O in water due to its high solubility of 0.0005855 mol m<sup>-3</sup> Pa<sup>-1</sup> at 0°C in fresh water (Weiss and Price, 1980), which implies that in equilibrium about 30% of the N<sub>2</sub>O is dissolved in the water. The equi-

- <sup>15</sup> librium isotope effect between N<sub>2</sub>O dissolved in water and N<sub>2</sub>O in the gas phase is known, -0.75% for <sup>15</sup>N/<sup>14</sup>N and -1.06% for <sup>18</sup>O/<sup>16</sup>O (Inoue and Mook, 1994), respectively. This would cause under equilibrium conditions a change of the N<sub>2</sub>O isotopic composition in the gas phase of less than 0.3‰ and 0.5‰ for  $\delta^{15}$ N and  $\delta^{18}$ O, respectively. However, during the entire extraction the gas phase above the water is constantly
- <sup>20</sup> pumped off, so that N<sub>2</sub>O is continuously removed, and we do not expect a significant isotope fractionation due to remaining dissolved N<sub>2</sub>O for our measurements. Also the kinetic fractionation, which would affect the results on incomplete extraction of N<sub>2</sub>O, is small,  $-0.7\pm\%$  for  $\delta^{15}$ N and -1.9% for  $\delta^{18}$ O (Inoue and Mook, 1994), and should thus have no significant effect on our results.
- The N<sub>2</sub>O isotopic composition (<sup>18</sup>O, average <sup>15</sup>N and position dependent <sup>15</sup>N) was measured using continuous-flow isotope ratio mass spectrometric techniques. The principle is described in Röckmann et al. (2003a, b). We only give a brief summary here. About 300 cm<sup>3</sup> of firn air are injected into a He carrier gas stream of a modified ThermoFinnigan PreCon system. After removal of most of CO<sub>2</sub> over Ascarite

7551

(NaOH coated silica), N<sub>2</sub>O is cryogenically preconcentrated. Final separation from the remaining CO<sub>2</sub> occurs on a capillary GC column (PoraPlot Q, 0.32 mm i.d., 25 m) after cryo-focusing the sample. The column is separated into a pre-column and an analytical column. The pre-column is backflushed after each measurement which eliminates

interferences from other atmospheric compounds, in particular on m/z 31 for quantifying the NO<sup>+</sup> fragment. The sample is then transferred to the IRMS via an open split interface.

For the measurements presented here, two different mass spectrometers were used. The firn air samples were analyzed on a ThermoFinnigan Delta plus XL and the ice air

- <sup>10</sup> samples on a ThermoFinnigan Delta plus XP. On the Delta plus XL, three collectors can be monitored simultaneously, therefore requiring two separate sample injections: one for the N<sub>2</sub>O<sup>+</sup> ions at m/z 44, 45 and 46, and one for the NO<sup>+</sup> fragment at m/z 30 and 31 (Brenninkmeijer and Röckmann, 1999; Röckmann et al., 2003b). On the Delta plus XP, all five mass-to-charge ratios can be monitored simultaneously. Thus,
- <sup>15</sup> only a single injection is needed (Röckmann and Levin, 2005). As laboratory reference gas, we use a carefully calibrated atmospheric air sample (Kaiser et al., 2003a) with a N<sub>2</sub>O mixing ratio of 319.0 ppbv and isotope ratios of  $\delta^{15}$ N=6.6‰ vs. atmospheric N<sub>2</sub> and  $\delta^{18}$ O=44.6‰ vs. VSMOW. The mass spectrometer reference gas (N<sub>2</sub>O, purity 99.9999%) has an isotopic composition of  $\delta^{15}$ N=0.53‰ vs. atmospheric N<sub>2</sub> and
- $^{20}$   $\delta^{18}$ O=39.60‰ vs. VSMOW. The calibration of the intramolecular distribution in N<sub>2</sub>O is not straightforward; a mass spectrometric calibration method using mixtures of isotopically labeled N<sub>2</sub>O isotopologues has recently been developed (Kaiser et al., 2004). This calibration will be used in this work, but we note that there exists a large difference to the only other calibration available (Toyoda and Yoshida, 1999). Using that calibra-
- tion shifts  ${}^{1}\delta^{15}$ N values up by about 21‰ and  ${}^{2}\delta^{15}$ N values down by about 21‰, so it is very important to compare the reference scale when comparing absolute  ${}^{1}\delta^{15}$ N and  ${}^{2}\delta^{15}$ N values from different publications. To distinguish between the N<sub>2</sub>O isotopomers,

we adopt the following notation:

<sup>15</sup>N terminal position <sup>15</sup>N<sup>14</sup>NO <sup>1</sup> $\delta$ <sup>15</sup>N

<sup>15</sup>N central position <sup>14</sup>N<sup>15</sup>NO  $^{2}\delta^{15}N$ 

The samples were measured three times and the standard deviations on the measured values in the firn are 0.05‰, 0.3‰, 0.3‰ and 0.1‰ for  $\delta^{15}$ N,  ${}^{1}\delta^{15}$ N,  ${}^{2}\delta^{15}$ N and  $\delta^{18}$ O, respectively. Due to the limited sample amount the ice core samples were only measured once, and the sample size was at most 100 cm<sup>3</sup>. The standard deviations are deduced from the tests with reference gas and bubble-free ice and amount to 0.4‰, 0.8‰, 0.8‰ and 0.5‰ for  $\delta^{15}$ N,  ${}^{1}\delta^{15}$ N,  ${}^{2}\delta^{15}$ N and  $\delta^{18}$ O, respectively.

## 10 3 Results

 $N_2O$  concentration and isotope profiles in firn and ice air are shown in Fig. 1. As expected from its atmospheric evolution, the  $N_2O$  mixing ratio at each sampling site decreases with depth. The overall shape of the  $N_2O$  isotopic signals is similar for both sites and we identify three zones with similar characteristics: (1) a small increase of

- the heavy isotopes from the surface to the depth where closed porosity (see description of the firn column in the next section) appears (50 to 65 m depending on the site);
  (2) a large increase from this particular depth down to the bottom part of the firn; (3) a stabilization or slower increase (but with large scatter of the measurements) within the ice. In Berkner Island, the gas extracted at 75 m exhibits depleted isotopic values.
- <sup>20</sup> ues for  $\delta^{15}$ N and  ${}^{1}\delta^{15}$ N. This is due to an unusually depleted measurement for  $\delta^{15}$ N, which affects strongly the  ${}^{15}$ N value calculated for the terminal position, whereas the central position, which is measured independently, is much less affected. Since no repeated measurements are possible on the ice air samples, this discrepancy cannot be explained. However, it is highly unlikely that this is indeed an atmospheric signal.
- <sup>25</sup> Therefore, we present this point in Fig. 1, but have excluded it from the further analysis. In addition, at this site, analyses conducted on other gases, like CH<sub>4</sub> and CFCs,

7553

showed that the firn air sampled between 58 m and the close-off depth (64 m) may be contaminated by ambient air. However, the  $CO_2$  mixing ratio was monitored using an infra-red analyzer (LICOR), during firn air sampling. Below 58 m, the  $CO_2$  mixing ratio was stable and below the ambient air level, thus contradicting the hypothesis of a contamination. As the validity of the last points is questionable, we decided to remove

contamination. As the validity of the last points is questionable, we decided to remove them and we consider the firn profile from the surface to 57 m. This observation does not concern the ice samples.

## 4 Modelling

## 4.1 Firn air model

- <sup>10</sup> Physical processes, like diffusion or gravitation, affect the trace-gas isotopic composition within the firn column; therefore, firn air measurements cannot be directly used to derive the atmospheric history of trace-gas isotopes. Firn air diffusion models take these effects into account and allow the selection of atmospheric scenarios compatible with the firn profile. In addition, N<sub>2</sub>O extracted from our ice samples dates from 1740
- <sup>15</sup> AD to 1910 AD, a time period when the atmospheric N<sub>2</sub>O mixing ratio increased. This increase could affect the air isotopic profile through diffusion along the concentration gradient. To take this potential effect into account, the composition of bubbly ice is also computed using the diffusion model.
- We employed a 1-D gas diffusion model in Eulerian coordinates developed by Rom-<sup>20</sup> melaere et al. (1997). The processes taken into account include (1) air mixing by pressure and temperature gradient down to a few meters from the surface (i.e. the so-called convection zone), (2) molecular diffusion in the open pore space and gravitational fractionation (entrainment toward the deeper firn depends on concentration gradients, diffusivities and molar mass), (3) a downward air flux in the open porosity
- <sup>25</sup> zone due to bubble closure removing air from open pores; this removed air has to be replaced by air coming from the upper parts of the firn, thus creating this downward

flux. These phenomena also affect the isotopologues of trace gases. The model computes them as independent species and calculates the resulting isotopic ratios in the firn afterwards. Temperature, accumulation rate and firn structural parameters (density, open and closed porosity, tortuosity) are set to their present-day measured values.

The diffusivity profile for any trace gas and the firn-ice transition depth thus remain constant with time. Assuming a constant temperature in our model, we are not resolving variations over the top 10 m, where seasonal temperature effects are imprinted.

One big "unknown" of the firn air model is the diffusivity profile at a specific location. Diffusivities have not been measured at the locations discussed here. We cal-

- culate the diffusivity profile from the well-known  $CO_2$  evolution in the atmosphere and our measured  $CO_2$  mixing ratio profile. As  $CO_2$  and  $N_2O$  both exhibit an approximately exponential increase since pre-industrial time,  $N_2O$  diffusivities are then calculated according to  $D(N_2O)/D(CO_2)=1.004$  (Trudinger et al., 1997). In order to validate the diffusivity profile, we apply the firn model using a historical record of  $N_2O$  mix-
- <sup>15</sup> ing ratio. This scenario is derived from an interpolation of ice core data (Flückiger et al., 1999), the scenario deduced from firn air measurements at South Pole (Battle et al., 1996), and direct atmospheric observations at Cape Grim, Tasmania, from the ALE/GAGE/AGAGE measurements (http://cdiac.esd.ornl.gov/ftp/ale\_gage\_Agage/ AGAGE/gc-md/complete/tasmania). The agreement between the N<sub>2</sub>O data and the <sup>20</sup> model reconstruction is good, as seen in Fig. 1, thus validating the density and diffu-
- sivity profiles used in the model.
  - 4.2 Monte-Carlo modelling of N<sub>2</sub>O evolution

Although routinely used for trace gas mixing ratios, the implementation of the inversion technique to isotope ratios in the firn air model is complicated, owing to the nonlinearity of the forward model in the case of isotopes. We use a Monte Carlo approach instead where a wide range of hypothetical scenarios for the atmospheric evolutions of isotopologue concentrations are tested using the firn air model in the forward mode. As the evolution of N<sub>2</sub>O mixing ratio since the pre-industrial era is close to an expo-

nential increase, we parameterize the input scenarios for the atmospheric  $\delta$  value as an exponential function. This is a good mathematical approximation for the assumption that one source with constant isotopic composition has been responsible for the change in isotopic composition since pre-industrial times. We note that this is a strong

- <sup>5</sup> constraint from the model, which does not allow that various sources with distinct isotopic composition have contributed differently at different times. Furthermore, we note already here that the fact that the atmospheric  $\delta$  value decreases does not necessarily mean that the isotopic composition of the sources also decreases. In a disequilibrium situation between sources and sinks, which is driven by strong source changes, the at-
- <sup>10</sup> mospheric  $\delta$  value will always change in the direction of the isotopic composition of the global average source. Since the global average N<sub>2</sub>O source is isotopically depleted versus atmospheric N<sub>2</sub>O, the atmospheric  $\delta$  value would also decrease if the source signature had stayed constant and even if it slightly increased.
- The validity of each scenario is assessed based on agreement between the mod-<sup>15</sup> elled and experimental firn profiles using a  $\chi^2$  test. A scenario is classified acceptable if the corresponding calculated isotope profile in the firn falls within a given range of the experimental data (Table 2) (Aballain, 2002; Bräunlich et al., 2001) The accepted scenarios form the envelope that is presented in the results. A measured atmospheric isotopic evolution was used as input data, from 1990 AD to present. This profile comes
- from direct atmospheric measurements on air samples collected in Neumayer, Antarctica (70° S, 8° W) from 1990 AD to 2002 AD (Röckmann and Levin, 2005). A trend on each isotopologue was deduced from this profile at this specific site. Then, starting from the measured surface value, this trend was set as a "known" isotopic evolution for the last 12 years at each studied site (i.e. NGRIP and Berkner Island), i.e. the isotopic
- evolution since 1990 was fixed. Note that the surface firn air values differ slightly between NGRIP and Berkner, but this difference is within the analytical uncertainty and does not alter the general trends derived from the firn air diffusion model.

<sup>7555</sup> 

#### 4.3 Comparison of model and data

For NGRIP, the model can reproduce the firn and ice data quite well, especially for  $\delta^{15}$ N and  $\delta^{18}$ O. The average difference between the model and the data is less than 0.15‰ for  $\delta^{15}$ N and less than 0.20‰ for  $\delta^{18}$ O. The input scenario, constrained by

- <sup>5</sup> direct isotope measurements between 1990 AD and the present, allows reproducing the top part of the firn within uncertainties. The  ${}^{1}\delta^{15}N$  and  ${}^{2}\delta^{15}N$  profiles are more scattered due to the difficulty of measuring the ( ${}^{15}N$ )N<sub>2</sub>O isotopomers. In addition to the generally larger scatter, the ice core results do not seem to extend the firn air data smoothly; for the  ${}^{1}\delta^{15}N$  the ice core data appear rather depleted, whereas they are
- rather enriched for  ${}^{2}\delta^{15}N$  compared to the firn data. Although the model is able to reproduce on average the trends of the data, the resulting scenarios should be regarded with caution, due to the larger uncertainty of the measurements on ice samples. Given the large experimental uncertainty of the ice core measurements and the fact that we only have two data points from ice air, we believe that the constraints from the ice data
- <sup>15</sup> are probably over-represented in our model. If we use the ice core data at their respective 1 $\sigma$  error limits, the ice and firn data are much more consistent. This indicates that the trends estimated from our combined data are likely underestimated for  ${}^{1}\delta^{15}N$  and overestimated for  ${}^{2}\delta^{15}N$ .
- Also in the case of Berkner Island, the model reproduces the measured firn profile well. The ice data are more scattered but taking into account the error bars, the data lie within the envelope of accepted scenarios defined by the  $\chi^2$  test, except for the point at 75 m which is very low for  $\delta^{15}$ N and  ${}^{1}\delta^{15}$ N (previously mentioned). As for NGRIP, the larger scatter of intramolecular  ${}^{15}$ N measurements implies more caution for the interpretation of the deduced atmospheric scenario.
- The reconstructed atmospheric trends for both sites and for each isotopologue are represented in Fig. 2, together with the experimental values of the ice samples corrected for gravitation. The colour scale represents the probability that a given atmospheric scenario is constrained by the firn data. The N<sub>2</sub>O in an air sample at a certain

7557

depth has a wide age spectrum due to mixing processes in the firn column. This age spectrum widens with depth, and in particular  $N_2O$  near the bottom of the firn is not representative for a certain atmospheric age. Thus, the constraints from the firn air measurements on the atmospheric scenarios become successively weaker with depth,

- and the probability that the derived atmospheric scenario is constrained by the firn data can be computed from the green functions deduced from the inverse model (Rommelaere et al., 1997). This probability is only calculated for gases in the open porosity region. Thus, it has no significance for the air extracted from the ice. We evaluate our results for two age ranges: one from 1970 AD to present, which is the well-constrained
- period (constrained by the firn data only), and one using all data. The latter is then primarily constrained by the ice air. The values are summarized in Table 3. For NGRIP, the colour scale shows that the firn data can constrain the atmospheric

scenario at, at least 50% from 1970 AD on. From 1970 AD to 2001 AD, the evolution of the isotopic composition of atmospheric N<sub>2</sub>O is about -1.1%, -1.1%, -1.1%, and -0.5% for  $\delta^{15}$ N,  $^{1}\delta^{15}$ N,  $^{2}\delta^{15}$ N and  $\delta^{18}$ O, respectively. The long term evolution, given by the isotopic to a given by the gi

by the ice data, leads to an overall decrease of the signal since 1700 AD of –2.9‰, –2.3‰, –3.3‰ and –1.3‰ for  $\delta^{15}$ N,  $^{1}\delta^{15}$ N,  $^{2}\delta^{15}$ N and  $\delta^{18}$ O, respectively.

20

For Berkner Island, the color scale indicates that the atmospheric scenario is constrained by the firn data at, at least 80% from 1970 AD on. From 1970 to 2003 AD, the evolution on the isotopic composition of atmospheric N<sub>2</sub>O is about -1.0%, -1.3%,

- -0.9% and -0.7% for  $\delta^{15}$ N,  ${}^{1}\delta^{15}$ N,  ${}^{2}\delta^{15}$ N and  $\delta^{18}$ O, respectively. Since 1700 AD, the depletion of the isotopic signal is about -2.7%, -2.5%, -3.1% and -2.1% or  $\delta^{15}$ N,  ${}^{1}\delta^{15}$ N,  ${}^{2}\delta^{15}$ N and  $\delta^{18}$ O, respectively.
- The depletion observed for the oxygen isotopes since pre-industrial times is smaller than that observed for the nitrogen isotopes. Consistent atmospheric scenarios are deduced at both sites within the error limits for the nitrogen isotopes. For the  $\delta^{18}$ O scenarios, slightly different results are derived for the two sites. Since the NGRIP scenarios are only constrained by two data points, the data from Berkner Island are likely more reliable, and we extend the range of possible scenarios from the best scenario for

NGRIP to the upper limit for Berkner Island. In addition, Fig. 2 shows that for  $\delta^{15}$ N and  $\delta^{18}$ O the trends measured recently on the archived Antarctic air samples (Neumayer) are in very good agreement with our results from firn and ice air. On the other hand, this is obviously not the case for individual <sup>15</sup>N signatures, where the trends from the direct measurements are too high for  $1\delta^{15}$ N and far too low for  $2\delta^{15}$ N, which produces

the kink in the isotope scenarios in Fig. 2.

A similar discrepancy between the Neumayer observations and the earlier firn data from Dome C and Dronning Maud Land (Röckmann et al., 2003a) was noted in Röckmann and Levin (2005). On the other hand the data from the different firn air

- <sup>10</sup> studies show similar trends (see below). The combined evidence from the old (Dome C and DML) and new (this study) firn and ice air data now indicates that indeed the trends at the two positions are not so different. Trends in  ${}^{2}\delta^{15}N$  are slightly larger than those for  ${}^{1}\delta^{15}N$  when the whole period is considered, but smaller in the recent period since 1970 AD. However, due the large uncertainties on the intramolecular measure-
- <sup>15</sup> ments, this is not significant.

## 5 Discussion

Only one previous study of  $N_2O$  isotopologues in pre-industrial ice has been performed so far, using ice from GISPII (Greenland) (Sowers et al., 2002). Sowers et al. found that the <sup>18</sup>O trends were not consistent between firn air and ice air measurements. In

- <sup>20</sup> particular, the isotope trends deduced from the ice core samples are much stronger. Note, however, that the analytical errors for ice core measurements (also in our measurements) are large. To explain this observation, the authors suggested a decoupling of the  $\delta^{15}$ N and  $\delta^{18}$ O trends over the last 200 years: the majority of  $\delta^{18}$ O change would have occurred in the 19th century whereas most of the  $\delta^{15}$ N change took place
- <sup>25</sup> during the 20th century. This would be isotopic evidence showing that the increase of N<sub>2</sub>O since pre-industrial times was not due to one common source with constant isotopic composition. However, in our study we deduce from both NGRIP and Berkner 7550

7559

Island firn and ice air samples consistent trends between the ice core and firn air results for  $\delta^{15}$ N and  $\delta^{18}$ O, which show a larger depletion of  $\delta^{15}$ N compared to  $\delta^{18}$ O, since 1700 AD. This does not confirm the proposed decoupling of  $\delta^{15}$ N and  $\delta^{18}$ O but rather suggests that they have changed in parallel over the entire time period. We can-

- <sup>5</sup> not completely rule out that our extraction method may introduce a  $\delta^{18}$ O enrichment of ~+0.5‰ (see Table 1). As mentioned above, we did not correct for this enrichment, which is at the limit of significance. However, accounting for this effect would only increase the difference between our  $\delta^{18}$ O results and the results of Sowers et al. (2002). Another possible explanation for the difference between both studies could lie in the
- <sup>10</sup> fractionation related to the N<sub>2</sub>O mixing ratio gradient between the atmosphere and the pore close-off during the early part of the atmospheric N<sub>2</sub>O increase. It amounted to ~0.02 ppbv yr<sup>-1</sup> between 1700 AD and 1900 AD. Sowers et al. (2002) only corrected their results for gravitational fractionation. On the other hand our model corrects for gravitation as well as for diffusion due to mixing ratio changes that have occurred since
- <sup>15</sup> 1700 AD. To check this hypothesis, we used the diffusion model to simulate  $\delta^{15}N$  at NGRIP in the forward mode. The input scenario included the historical [N<sub>2</sub>O] record and the atmospheric  $\delta^{15}N(N_2O)$  was kept constant over time. In this case the resulting signal in the ice has an amplitude of 0.32‰. The gravitational fractionation in NGRIP has been measured (A. Landais, personal communication) and is close to 0.32‰ for
- $\delta^{15}$ N. We deduce that the increase of N<sub>2</sub>O mixing ratio between 1700 AD and 1900 AD is not sharp enough to introduce a measurable fractionation; thus, the different model corrections applied by Sowers et al. (2002) and our analysis cannot explain the difference between the two sets of results.

Finally, it appears that our measurements have smaller errors, and that the larger errors in the Sowers et al. (2002) study provide only rather weak constraints that such a decoupling has actually taken place. We can thus conclude that the observed  $\delta^{18}$ O trend is smaller than the  $\delta^{15}$ N trend since pre-industrial times.

The  $\delta^{18}$ O of N<sub>2</sub>O reflects the oxygen isotopic signatures of its different sources such as molecular oxygen, soil water (nitrification) and substrate NO<sub>3</sub><sup>-</sup> (denitrification). It also

reflects the oxygen isotope enrichment factors associated with each reaction step. So far, it has not been possible yet to determine the relative contribution of these different oxygen sources in today's  $N_2O$  budget, because no consistent picture has yet emerged (Gros et al., 2004). Recent studies have also emphasized the importance of the population of microorganisms involved in the <sup>18</sup>O-enrichment (Schmidt et al., 2004; Sutka

et al., 2003).

In addition, since N<sub>2</sub>O sources and sinks are presently not in equilibrium, the isotopic composition of tropospheric N<sub>2</sub>O is shifting from its pre-industrial value towards the isotopic composition of the global average sources. The total offset between the

- global average source and the troposphere is determined by the fractionations in the stratosphere, and these are smaller for <sup>18</sup>O than for <sup>15</sup>N due to the smaller stratospheric fractionation constants (Kaiser et al., 2002b, 2003b; Röckmann et al., 2001a). When the atmosphere is moving towards the source isotopic composition, the absolute shifts are smaller for <sup>18</sup>O than for <sup>15</sup>N, simply because the isotopic difference to the global average sources is smaller for δ<sup>18</sup>O.
- Our results show that  $\delta^{18}$ O and  $\delta^{15}$ N have both obviously decreased since 1700 AD. Making the assumptions that the N<sub>2</sub>O lifetime, the stratospheric fractionation constants and the average troposphere/stratosphere exchange rate have remained constant since 1700 AD, Röckmann et al. (2003a) suggested that the depletion of the tro-
- <sup>20</sup> pospheric signal is due to an increase of sources depleted in heavy isotopes, whereas the flux of isotopically enriched N<sub>2</sub>O from the stratosphere to the troposphere has not had time to increase proportionally yet. As natural sources have probably remained constant, the source responsible for the depletion may thus be a depleted anthropogenic source. Agriculture represents a major depleted anthropogenic source and
- its contribution to the N<sub>2</sub>O global budget has increased since 1900 AD (Kroeze et al., 1999). The intensification of agriculture, at first order proportional to the population increase, involved an intense use of natural fertilizers until the middle of the 20th century, followed by the enhanced use of synthetic fertilizers after 1930 AD. Perez et al. (2000, 2001) suggested that synthetic fertilizers may be more depleted in heavy isotopes than

7561

natural ones, implying that recent agricultural-derived  $N_2O$  emissions are even more depleted than about one century ago. Assuming that half of the increased  $N_2O$  emission from agriculture during the past century has been from application of synthetic fertilizers, whereas the other half has come from natural fertilizers and indirect emission.

- sions (nitrate leaching and runoff), Pérez et al. (2001) predicts a decrease in  $\delta^{15}$ N of tropospheric N<sub>2</sub>O since pre-industrial times by 2 to 3‰This deduction is in good agreement with our results. Therefore, the decrease of the isotope ratios, observed for all isotopologues, could be explained by the increase of anthropogenic sources, agriculture in particular.
- <sup>10</sup> Röckmann et al. (2003a) performed measurements of the <sup>18</sup>O and positiondependent <sup>15</sup>N isotopic composition of N<sub>2</sub>O from Antarctic firn air sampled at Dome C (75° S, 123° W) and Dronning Maud Land (77° S, 10° W). By comparing these firn data to simulations carried out with the same firn air diffusion model as ours (without using the Monte-Carlo model), they reconstructed the temporal evolution of the
- $^{15}$  N<sub>2</sub>O isotope signatures, since pre-industrial times. However, the long-term trend that they reported is not constrained by ice air values since no ice samples were analyzed. Nevertheless, the trends deduced in the two studies from the firn data, with different approaches are quite consistent. By using the Green function, the age of N<sub>2</sub>O at the firn-ice transition, as well as its distribution has been calculated for all sites. The mean
- age of N<sub>2</sub>O is about 35, 40, 60 and 90 years old at Dome C, Dronning Maud Land, NGRIP and Berkner Island, respectively. Note, however, that in particular near the firn-ice transition, the age spectrum has a very wide distribution. So, for all sites, the evolution of the atmospheric isotopic signal can be constrained by the data measured in the firn over the last 30 years, i.e. until 1970 AD. Those trends are summarized in
- <sup>25</sup> Table 4. The atmospheric histories of the isotope ratios derived in both studies are similar for all isotopologues. There is a slight difference in the position-dependent <sup>15</sup>N values, but this is within the combined error range. In general, consistent results are found for all locations. They show that <sup>15</sup>N depletion is distributed quite evenly within the N<sub>2</sub>O molecule.

At this time, it is still difficult to draw conclusions from our intramolecular nitrogen results due to the high experimental errors as discussed above. Nevertheless, several tracks could be followed. Previous studies have shown that the stratospheric <sup>15</sup>N enrichments are position-dependent (Griffith et al., 2000; Park et al., 2004; Röckmann et al., 2001a; Toyoda et al., 2004). Whereas <sup>15</sup>N<sup>14</sup>NO is photolyzed faster than <sup>14</sup>N<sup>15</sup>NO,

- <sup>5</sup> al., 2001a; Toyoda et al., 2004). Whereas <sup>15</sup>N<sup>14</sup>NO is photolyzed faster than <sup>14</sup>N<sup>15</sup>NO, <sup>14</sup>N<sup>15</sup>NO is removed faster in the reaction with O(<sup>1</sup>D) (Hessberg et al., 2004; Kaiser et al., 2002a, 2003b; Röckmann et al., 2001a). This difference provides the possibility to distinguish between the two processes in the atmosphere based on the isotope fractionation pattern they leave in N<sub>2</sub>O. In principle this may allow to investigate both N<sub>2</sub>O
- stratospheric sinks since the pre-industrial era, but this has to be distinguished from variations in the two main production processes. It seems that nitrification, favored in dry and aerobic systems, preferentially enriches the central N position, compared to denitrification, favored in wet and anaerobic systems (Pérez et al., 2001). The enzymatic reaction pathways involved in N<sub>2</sub>O production may also suggest that the site
- <sup>15</sup> preference could be characteristic from the involved microorganisms. The intramolecular distribution of <sup>15</sup>N in N<sub>2</sub>O could thus provide a basis for distinguishing biological processes (Schmidt et al., 2004; Sutka et al., 2003). Further work needs to be performed to improve the precision of N position dependent measurements, so that they could add another constraint on the N<sub>2</sub>O evolution.

## 20 6 Conclusions

Isotope measurements on N<sub>2</sub>O from both Antarctic and Arctic firn and ice air samples show that the observed increase in the atmosphere since pre-industrial times is accompanied by a significant isotopic depletion in <sup>15</sup>N at both positions in the molecule and in <sup>18</sup>O. Using the data in combination with a firn air diffusion model, the evolution

<sup>25</sup> of the isotope signatures has been precisely quantified, which is important for global isotope budget calculations.

Consistent trends are found for <sup>15</sup>N and <sup>18</sup>O changes derived from firn air since

1970, from ice air since pre-industrial times, and from recent direct measurements on Antarctic air samples dating back to 1990 (Röckmann and Levin, 2005). The trends for different time periods are less consistent for the <sup>15</sup>N-isotopomers. In particular, the trends derived from the recent air samples are much stronger for <sup>1</sup> $\delta$ <sup>15</sup>N than for <sup>2</sup> $\delta$ <sup>15</sup>N,

whereas the firn and ice core data imply that they are more similar. This is unlikely to reflect an atmospheric signal and may be due to presently not resolved systematic errors in the analytical procedure.

The depletion of the signal for all the isotopologues can be partly related to the increase of agriculture, in particular to the use of fertilizers and is in good agreement with

- predictions made using bottom-up calculations. The different isotope signatures may potentially be used to discriminate between source and sink effects as origin of variations in the N<sub>2</sub>O mixing ratio on long time scales. Further work needs to be performed to improve the precision on the intramolecular <sup>15</sup>N measurements, in order to reduce the uncertainty and to better quantify weak isotopic variations.
- Acknowledgements. This work is a contribution to the European Commission project CRYO-STAT (EVK2-CT2001-00116) funded under the Energy, Environment and Sustainable Development Programme; 1998–2002. Laboratory work was also funded by the German AFO2000 program within the project ISOSTRAT and by the French-German "Programme d'Actions Intégrées" PROCOPE. The NorthGRIP project is directed and organized by the Department
- of Geophysics at the Niels Bohr Institute for Astronomy, Physics and Geophysics, University of Copenhagen. It is being supported by Funding Agencies in Denmark (SNF), Belgium (FNRS-CFB), France (IPEV and INSU/CNRS), Germany (AWI), Iceland (RannIs), Japan (MEXT), Sweden (SPRS), Switzerland (SNF) and the United States of America (NSF). The Berkner Island field project is a collaborative programme supported by the British Antarctic Survey, the french
- IPEV and INSU/CNRS. We thank all participants to the field work at both sites for drilling and firn air sampling operations. We also thank B. Knape for help with the isotope measurements in the laboratory.

<sup>7563</sup> 

#### References

30

- Aballain, O.: Reconstruction de l'évolution passée du rapport isotopique <sup>13</sup>C/<sup>12</sup>C du méthane atmosphérique, à partir de l'analyse de l'air extrait du névé polaire, Université Joseph Fourier Grenoble I, Grenoble, 2002.
- Battle, M., Bender, M., Sowers, T., Tans, P. P., Butler, J. H., Elkins, J. W., Ellis, J. T., Conway, T., Zhang, N., Lang, P., and Clarke, A. D.: Atmospheric gas concentrations over the past century measured in air from firn at the South Pole, Nature, 383, 231–235, 1996.
- Bräunlich, M., Aballain, O., Marik, T., Jockel, P., Brenninkmeijer, C. A. M., Chappellaz, J., Barnola, J. M., Mulvaney, R., and Sturges, W. T.: Changes in the global atmospheric
- methane budget over the last decades inferred from C-13 and D isotopic analysis of Antarctic firn air, J. Geophys. Res., 106, 20465–20481, 2001.
  - Brenninkmeijer, C. A. M. and Röckmann, T.: Mass spectrometry of the intramolecular nitrogen isotope distribution of environmental nitrous oxide using fragment-ion analysis, Rap. Commun. Mass Spectrom., 13, 2028–2033, 1999.
- <sup>15</sup> Crutzen, P. J.: The influence of nitrogen oxides on the atmospheric ozone content, Quart. J. R. Met. Soc., 96, 320–325, 1970.
  - Dore, J. E., Popp, B. N., Karl, D. M., and Sansone, F. J.: A large source of atmospheric nitrous oxide from subtropical North Pacific waters, Nature, 396, 33–66, 1998.
- Flückiger, J., Dallenbach, A., Blunier, T., Stauffer, B., Stocker, T. F., Raynaud, D., and Barnola,
   J. M.: Variations in atmospheric N<sub>2</sub>O concentration during abrupt climatic changes, Science, 285, 227–230, 1999.
  - Griffith, D. W. T., Toon, G. C., Sen, B., Blavier, J.-F., and Toth, R. A.: Vertical profiles of nitrous oxide isotopomer fractionation measured in the stratosphere, Geophys. Res. Lett., 27, 2485–2488, 2000.
- Gros, V., Brenninkmeijer, C. A. M., Jöckel, P., Kaiser, J., Lowry, D., Nisbet, E. G., O'Brian, P., Röckmann, T., and Warwick, N.: Isotope signatures of trace gas sources, in: Emissions Of Atmospheric Trace Compounds, edited by: Granier, C., Artaxo, P., and Reeves, C. E., Kluwer Academic Pub., Paris, 2004.
  - Inoue, H. Y. and Mook, W. G.: Equilibrium and kinetic nitrogen and oxygen isotope fractionations between dissolved and gaseous N<sub>2</sub>O, Chem. Geol., 113, 135–148, 1994.
- IPCC: Climate change 2001, The scientific basis, Cambridge University Press, 2001. Kaiser, J., Brenninkmeijer, C. A. M., and Röckmann, T.: Intramolecular <sup>15</sup>N and <sup>18</sup>O fractiona-

7565

tion in the reaction of  $N_2O$  with  $O(^1D)$  and its implications for the stratospheric  $N_2O$  isotope signature, J. Geophys. Res., 107, doi:10.1029/2001JD001506, 2002a.

- Kaiser, J., Röckmann, T., and Brenninkmeijer, C. A. M.: Temperature dependence of isotope fractionation in N<sub>2</sub>O photolysis, Phys. Chem. Chem. Phys., 4, 4420–4430, doi:10.1039/b204837j, 2002b.
- Kaiser, J., Röckmann, T., and Brenninkmeijer, C. A. M.: Complete and accurate mass spectrometric isotope analysis of tropospheric nitrous oxide, J. Geophys. Res., 108, doi:10.1029/2003JD003613, 2003a.
- Kaiser, J., Röckmann, T., Brenninkmeijer, C. A. M., and Crutzen, P. J.: Wavelength dependence of isotope fractionation in N<sub>2</sub>O photolysis, Atmos. Chem. Phys., 3, 303–313, 2003b,

SRef-ID: 1680-7324/acp/2003-3-303.

- Kaiser, J., Park, S., Boering, K. A., Brenninkmeijer, C. A. M., Hilkert, A. W., and Röckmann, T.: Mass-spectrometric method for the absolute calibration of the intramolecular nitrogen isotope distribution in nitrous oxide, Anal. Bioanal. Chem., 256-269, doi:10.1007/s00216-003-2233-2, 2004.
- Kim, K.-R. and Craig, H.: Nitrogen-15 and oxygen-18 characteristics of nitrous oxide: A global perspective, Science, 262, 1855–1857, 1993.
  - Kroeze, C., Mosier, A., and Bouwman, L.: Closing the N<sub>2</sub>O budget: A retrospective analysis 1500–1994, Glob. Biogeochem. Cycl., 13, 1–8, 1999.
- Minschwaner, K., Salawitsch, R. J., and McElroy, M. B.: Absorption of solar radiation by O<sub>2</sub>: Implications for O<sub>3</sub> and lifetimes of N<sub>2</sub>O, CFCl<sub>3</sub>, and CF<sub>2</sub>Cl<sub>2</sub>, J. Geophys. Res., 98, 10543– 10561, 1993.
- Naqvi, S. W. A., Yoshinari, T., Jayakumar, D. A., Altabet, M. A., Narvekar, P. V., Devol, A. H., Brandes, J. A., and Codispoti, L. A.: Budgetary and biogeochemical implications of N<sub>2</sub>O isotope signatures in the Arabian Sea, Nature, 394, 462–464, 1998.
- Park, S. Y., Atlas, E. L., and Boering, K. A.: Measurements of N<sub>2</sub>O isotopologues in the stratosphere: Influence of transport on the apparent enrichment factors and the isotopologue fluxes to the troposphere, J. Geophys. Res., 109, doi:10.1029/2003JD003731, 2004.
- Pérez, T., Trumbore, S. E., Tyler, S. C., Davidson, E. A., Keller, M., and De Camargo, P. B.:
   Isotopic variability of N<sub>2</sub>O emissions from tropical forest soils, Glob. Biogeochem. Cycl., 14, 525–535, 2000.
  - Pérez, T., Trumbore, S. E., Tyler, S. C., Matson, P. A., Ortiz-Monasterio, I., Rahn, T., and Griffith, D. W. T.: Identifying the agricultural imprint on the global N<sub>2</sub>O budget using stable isotopes,

J. Geophys. Res., 106, 9869–9878, 2001.

10

- Rahn, T. and Wahlen, M.: A reassessment of the global isotopic budget of atmospheric nitrous oxide, Glob. Biogeochem. Cycl., 14, 537–543, 2000.
- Röckmann, T., Kaiser, J., and Brenninkmeijer, C. A. M.: The isotopic fingerprint of the preindustrial and the anthropogenic N<sub>2</sub>O source, Atmos. Chem. Phys., 3, 315–323, 2003a, SRef-ID: 1680-7324/acp/2003-3-315.
  - Röckmann, T., Kaiser, J., Brenninkmeijer, C. A. M., and Brand, W. A.: Gas chromatography/isotope-ratio mass spectrometry method for high-precision position-dependent <sup>15</sup>N and <sup>18</sup>O measurements of atmospheric nitrous oxide, Rap. Commun. Mass Spectrom., 17, 1897–1908, 2003b.
- Röckmann, T., Kaiser, J., Brenninkmeijer, C. A. M., Crowley, J. N., Borchers, R., Brand, W. A., and Crutzen, P. J.: Isotopic enrichment of nitrous oxide (<sup>15</sup>N<sup>14</sup>NO, <sup>14</sup>N<sup>15</sup>NO, <sup>14</sup>N<sup>14</sup>N<sup>18</sup>O) in the stratosphere and in the laboratory, J. Geophys. Res., 106, 10403–10410, 2001a.
- Röckmann, T., Kaiser, J., Crowley, J. N., Brenninkmeijer, C. A. M., and Crutzen, P. J.: The origin
   of the anomalous or "mass-independent" oxygen isotope fractionation in tropospheric N<sub>2</sub>O,
   Geophys. Res. Lett., 28, 503–506, 2001b.
  - Röckmann, T. and Levin, I.: High-precision determination of the changing isotopic composition of atmospheric  $N_2O$  from 1990 to 2002, J. Geophys. Res., in press, doi:10.1029/2005JD006066, 2005.
- 20 Rommelaere, V., Arnaud, L., and Barnola, J.-M.: Reconstructing recent atmospheric trace gas concentrations from polar firn and bubbly sea ice data by inverse methods, J. Geophys. Res., 102, 30 069–30 083, 1997.
  - Schmidt, H.-L., Werner, R. A., Yoshida, N., and Well, R.: Is the isotopic composition of nitrous oxide an indicator for its origin from nitrification or denitrification? A theoretical approach
- from referred data and microbiological and enzyme kinetic aspects, Rapid Commun. Mass. Spectrom., 18, 2036–2040, 2004.

Sowers, T., Alley, R. B., and Jubenville, J.: Ice core records of atmospheric N<sub>2</sub>O covering the last 106 000 years, Science, 301, 945–948, 2003.

- Sowers, T., Rodebaugh, A., Yoshida, N., and Toyoda, S.: Extending records of the iso-
- topic composition of atmospheric N<sub>2</sub>O back to 1800 AD from air trapped in snow at the South Pole and the Greenland Ice Sheet Project II ice core, Global Biogeochem. Cycl., 16, doi:10.1029/2002GB001911, 2002.

Sutka, R. L., Ostrom, N. E., Ostrom, P. H., Gandhi, H., and Breznak, J. A.: Nitrogen isotopomer

7567

site preference of  $N_2O$  produced by Nitrosomonas europaea and Methylococcus capsulatus Bath, Rapid Commun. Mass Spectrom., 17, 738–745, 2003.

- Toyoda, S. and Yoshida, N.: Determination of nitrogen isotopomers of nitrous oxide on a modified isotope ratio mass spectrometer, Anal. Chem., 71, 4711–4718, 1999.
- <sup>5</sup> Toyoda, S., Yoshida, N., Urabe, T., Nakayama, Y., Suzuki, T., Tsuji, K., Shibuya, K., Aoki, S., Nakazawa, T., Ishidoya, S., Ishijima, K., Sugawara, S., Machida, T., Hashida, G., Morimoto, S., and Honda, H.: Temporal and latitudinal distributions of stratospheric N<sub>2</sub>O isotopomers, J. Geophys. Res., 109, doi:10.1029/2003JD004316, 2004.
- Trudinger, C. M., Enting, D. M., Etheridge, D. M., Francey, R. J., Levchenko, V. A., Steele,
   L. P., Raynaud, D., and Arnaud, L.: Modeling air movement and bubble trapping in firn, J.
   Geophys. Res., 102, 6747–6763, 1997.

von Hessberg, P., Kaiser, J., Enghoff, M. B., McLinden, C. A., Sorensen, S. L., Röckmann, T., and Johnson, M. S.: Ultra-violet absorption cross sections of isotopically substituted nitrous oxide species: <sup>14</sup>N<sup>14</sup>NO, <sup>15</sup>N<sup>14</sup>NO, <sup>14</sup>N<sup>15</sup>NO and <sup>15</sup>N<sup>15</sup>NO, Atmos. Chem. Phys., 4, 1237– 1253, 2004.

SRef-ID: 1680-7324/acp/2004-4-1237.

15

20

Weiss, R. F. and Price, B. A.: Nitrous-Oxide Solubility in Water and Seawater, Marine Chem., 8, 347–359, 1980.

Yoshida, N. and Toyoda, S.: Constraining the atmospheric N<sub>2</sub>O budget from intramolecular site preference in N<sub>2</sub>O isotopomers, Nature, 405, 330–334, 2000.

		$^{1}\delta^{15}$ N/‰ (vs. atm. N <sub>2</sub> )	$^{2}\delta^{15}$ N/‰ (vs. atm. N <sub>2</sub> )	$\delta^{18}$ O/‰ (vs. VSMOW)
Stand SM25 + bubble free ice	6.6±0.1	-16.9±0.8	30.1±1.0	44.5±0.5
Stand SM25	6.72±0.01	-16.4±1.4	29.8±1.4	44.0±0.2

Table 1. Comparison between the resulting  $N_2O$  isotopic ratios after extraction and the standard gas.

Table 2. Parameters that are used in Monte Carlo runs of the firn air diffusion model to qualify	
calculated scenarios as acceptable or rejected.	

	$\delta^{15}$ N	$^{1}\delta^{15}$ N	$\delta^{15}$ N	$\delta^{18}$ O
Monte Carlo parameters (‰) NGRIP Berkner Island	±0.3 ±1	±0.5 ±2	±2 ±1.4	±0.3 ±0.9

	NGRIP (‰)	Range (‰)	Berkner Island (‰)	Range (‰)	Average value (‰)	Range (‰)
1700 AD						
$\delta^{15}$ N	-2.9	-3.0 to -2.7	-2.7	-3.4 to -1.8	-2.8	-3.4 to -1.8
$^{1}\delta^{15}N$	-2.3	-2.3 to -1.6	-2.5	-3.5 to -1.5	-2.4	-3.5 to -1.5
$^{2}\delta^{15}N$	-3.3	-3.3 to -2.2	-3.1	-4.0 to -2.2	-3.2	-4.0 to -2.2
$\delta^{18}$ O	-1.3	-1.3 to -0.9	-2.1	-2.5 to -1.7	-1.6	-2.5 to -1.3
1970 AD						
$\delta^{15}$ N	-1.1	-1.1 to -1.0	-1.0	-1.2 to -0.8	-1.1	-1.2 to -0.8
$^{1}\delta^{15}N$	-1.1	-1.2 to -1.0	-1.3	-1.5 to -1.2	-1.2	-1.5 to -1.0
$\delta^{15}$ N	-1.1	-1.1 to -0.7	-0.9	-1.2 to -0.7	-1.0	-1.2 to -0.7
$\delta^{18}$ O	-0.5	-0.5 to -0.4	-0.7	-0.8 to -0.6	-0.6	-0.8 to -0.5


**Table 3.** Atmospheric change of  $\delta^{15}$ N,  ${}^{1}\delta^{15}$ N,  ${}^{2}\delta^{15}$ N and  $\delta^{18}$ O since pre-industrial times in NGRIP, Berkner Island and average value.

7571


Table 4. Evolution of N<sub>2</sub>O isotopic signatures since 1970 AD in different polar sites.

	NGRIP / Berkner Island <sup>(a)</sup> (‰)	Range (‰)	Dome C / DML <sup>(b),(c)</sup> (‰)
0 11	-1.1	-1.2 to -0.8	-0.9
$^{1}\delta^{15}$ N	-1.2	-1.5 to -1.0	-0.8
$^{2}\delta^{15}$ N	-1.0	-1.2 to -0.7	-1.0
δ <sup>18</sup> Ο	-0.6	-0.8 to -0.5	-0.5

<sup>(a)</sup> this study
 <sup>(b)</sup> Röckmann et al. (2003a)
 <sup>(c)</sup> range not modelled



**Fig. 1.** Mixing and isotopic ratios of  $N_2O$  in NGRIP and Berkner Island. Square: firn samples; diamonds: ice samples; solid line: best scenario; dashed lines: accepted scenarios. In some cases the upper or lower envelope coincides with the best scenario.



**Fig. 2.** Evolution of the atmospheric isotopic composition of N<sub>2</sub>O since 1700 AD in NGRIP and Berkner Island. Dashed lines: envelope of accepted scenarios; solid line: best scenario; green diamonds: ice samples corrected for gravitational fractionation. In Berkner Island, for  ${}^{1}\delta^{15}N$  two data points are missing because they are off scale. The kinks in the scenarios for the individual  ${}^{15}N$  isotopomers (see continuation of figure on next page) indicate a discrepancy between the scenarios presented here and the direct measurements on Antarctic air samples since 1990 (Röckmann and Levin, 2005); see text for further discussion.

7573




Fig. 2. Continued.

7575